



EPILEPSY

Epilepsy is disorder of the interconnections of the brain, which means people sometimes have seizures. Epilepsy means you are likely to have more than one seizure. Not everyone who has a seizure has epilepsy. The mainstay of epilepsy treatment is medications however there is a small proportion whereby neurosurgical intervention will be beneficial.

CAUSE

People often wonder what causes epilepsy. In about 50 per cent of cases, doctors cannot find a cause - epilepsy just seems to come out of the blue. In the other 50 per cent of the cases, the cause can be identified, just like any injury or illness to the brain.

The structural lesions causing epilepsy include:

- Hippocampal or Mesial Temporal Sclerosis
- Tumours
- Glioma
- Dysembryoplastic Neuroepithelial Tumour
- Meningioma
- Cerebrovascular lesions
- Cavemous Angioma
- Arteriovenous Malformation
- Disorders of Neuronal Migration and Organisation
- Encephalomalacia

Epilepsy occurring after head injury is called post-traumatic epilepsy and may be a result of scarring within the brain substance.

TYPES OF SEIZURES

Tonic-Clonic seizure

This typically begins with a tonic phase which lasts around 10-15 seconds during which time the patient may loss consciousness and collapse. The patient's body stiffens, teeth clench shut and stop breathing. There may be urinary incontinence and tongue biting at this stage. The clonic phase then begins, characterized by rhythmical muscular contractions involving the whole body.

The entire seizure usually lasts one to three minutes. A post-ictal state may occur following the seizure whereby the patient is confused and drowsy for a short period of time.

Absence seizures

This type of seizure normally affects children and lasts only 5 to 10 seconds. There is a brief loss of consciousness but no collapse or motor twitches/contractions apart from rapid blinking of the eyes. The patient normally returns to normal abruptly with no recollection of the event following the seizure.

Partial seizures

These may be simple or complex. Simple partial seizures do not involve changes in conscious state whilst complex partial seizures usually demonstrate impaired conscious states. Complex partial seizures may also begin with an aura such as fear, unusual smells or tastes or a feeling of déjà vu.

Partial seizure may progress to a tonic-clonic seizure called a secondarily generalized tonic-clonic seizure.

INVESTIGATIONS

The diagnosis of epilepsy is through a thorough history and examination of the patient. An eye-witness account of seizures is important in characterizing the type of seizure and to this end video monitoring may be used. The pre-operative work-up of an epilepsy surgery patient is complex and involved including several of the following.

- Blood tests – there are no specific blood tests to diagnose epilepsy. An FBE, electrolytes and coagulation profile are needed prior to any neurosurgical intervention. Regular drug level checks of anti-epileptic medications will be performed by the neurologist.
- Electroencephalogram (EEG) – this is performed to monitor the electrical interconnections of the brain. It is able to localize abnormal electrical or seizure activity and is used in the immediate post-ictal and inter-ictal period. Video EEG monitoring may be performed in hospital for one day up to one week with seizure stimuli such as sleep deprivation.
- Radiological tests
 - CT brain – this is performed to rule out the presence of a structural lesion as the cause of the seizure (eg tumour, vascular malformation)
 - MRI brain – this is performed to further delineate any underlying brain abnormalities and also gives information with regards to brain volumes
 - SPECT/PET scans – these are performed in the pre-surgical work-up to localize the specific region of abnormality and side of pathology.
- Invasive EEG monitoring – this includes intra-cranial EEG monitoring with subdural grids and bilateral depth electrodes. Further discussion of these are found under operations – Epilepsy surgery.

What type of management is required for people with epilepsy?

Suitable control of seizures with anti-convulsants (medication). Understanding their type of epilepsy, the warning signs, triggering factors and a change of life style to accommodate their condition. Surgical options for the treatment of patients with epilepsy not controlled by medication is increasingly successful. Suitable candidates may be offered surgery.

St Vincent's Private Hospital Melbourne

St Vincent's Private Hospital Fitzroy
Phone: (03) 9411 7111

Website: www.svphm.org.au

St Vincent's Private Hospital East Melbourne
Phone: (03) 9928 6555

Website: www.svphm.org.au

St Vincent's Hospital Melbourne

St Vincent's Hospital Fitzroy
Telephone: (03) 9231 2211

Website: www.svhm.org.au

Neurosurgery

Dr. Kristian Bulluss
Phone: (03) 9416 4619

Dr. Peter McNeill
Phone: (03) 9928 6333

Dr. Paul Smith
Phone: (03) 9639 3889

Dr. Carlos Chung
Phone: (03) 9419 5597

Assoc. Prof. Michael Murphy
Phone: (03) 9416 4619

Dr. Christopher Thien
Phone: (03) 9421 0355

Dr. Tiew Han
Phone: (03) 03 9417

Dr. Brendan O'Brien
Phone: (03) 9417 5033

Dr. Yi Yuen (Ian) Wang
Phone: (03) 9939 7112

Neurology

Prof. Mark Cook
Phone: (03) 9288 3068